



## *A Priori* Bounds for Periodic Solutions of a Delay Rayleigh Equation

GEN-QIANG WANG

Department of Mathematics, Hanshan Teacher's College  
Chaozhou, Guangdong 521041, P.R. China

SUI SUN CHENG

Department of Mathematics, Tsing Hua University  
Hsinchu, Taiwan 30043, R.O.C.

(Received and accepted April 1998)

**Abstract**—*A priori* bounds are established for periodic solutions of a Rayleigh equation with delay. By means of these bounds, an existence theorem for periodic solutions can be obtained by means of Mawhin's continuation theorem. © 1999 Elsevier Science Ltd. All rights reserved.

**Keywords**—Rayleigh equation, Delay, Periodic solution.

In [1], continuation theorems are introduced and applied to the existence of solutions of differential equations. In particular, a specific example is given in [1, p. 99] (see also [2, p. 175]) on how periodic solutions can be obtained by means of these theorems for the differential equation

$$x''(t) + f(x'(t)) + h(t, x(t)) = 0,$$

where  $f$  is a real continuous function defined on  $R$  such that  $f(0) = 0$ ,  $h$  is a real continuous function defined on  $R \times R$ ,  $2\pi$ -periodic in  $t$  and  $h(t, x)x < 0$  for  $|x| \geq r$  and  $t \in [0, 2\pi]$ . In the course of derivations, it is realized that once appropriate *a priori* bounds for the  $2\pi$ -periodic solutions of the equation

$$x''(t) + \lambda f(x'(t)) + \lambda h(t, x(t)) = 0$$

are known for each  $\lambda \in (0, 1)$ , then standard procedures will allow these theorems to imply existence of periodic solutions to the original equation.

In this note, we will be concerned with a similar equation with an additional delay

$$x''(t) + \lambda f(x'(t)) + \lambda g(x(t - \tau(t))) = \lambda p(t), \quad \lambda \in (0, 1), \quad (1)$$

where  $f$ ,  $g$ ,  $p$ , and  $\tau$  are real continuous functions defined on  $R$  such that  $f(0) = 0$ ,  $\tau$  and  $p$  are periodic with period  $2\pi$ , and

$$\int_0^{2\pi} p(t) dt = 0.$$

We will establish *a priori* bounds for solutions of this equation under relatively simple conditions on  $f$  and  $g$ . Once these bounds are obtained, existence of periodic solutions can be demonstrated for the equation

$$x''(t) + f(x'(t)) + g(x(t - \tau(t))) = p(t). \quad (2)$$

We remark that there are a number of studies which are concerned with the existence of periodic solutions of differential equations by means of establishing *a priori* bounds (see, e.g., [3]). Other related results can be found in [4–6].

**THEOREM 1.** *Suppose there are positive constants  $K$ ,  $D$ , and  $M$  such that*

- (i)  $|f(x)| \leq K$  for  $x \in R$ ,
- (ii)  $xg(x) > 0$  and  $|g(x)| > K$  for  $|x| \geq D$ , and
- (iii)  $g(x) \geq -M$  for  $x \leq -D$ .

*Then for any  $2\pi$ -periodic solution  $x = x(t)$  of (1),  $|x'(t)| \leq D_2$  and  $|x(t)| \leq D + 2\pi D_2$  for some positive number  $D_2$ .*

**PROOF.** Let  $x = x(t)$  be a  $2\pi$ -periodic solution of the equation (1). Since  $x(0) = x(2\pi)$ , there is some  $t_0 \in [0, 2\pi]$  such that  $x'(t_0) = 0$ . In view of (1), we see that for any  $t \in [0, 2\pi]$ ,

$$\begin{aligned} |x'(t)| &= \left| \int_{t_0}^t x''(s) ds \right| \leq \int_0^{2\pi} |x''(s)| ds \\ &\leq \lambda \int_0^{2\pi} |f(x'(s))| ds + \lambda \int_0^{2\pi} |g(x(s - \tau(s)))| ds + \lambda \int_0^{2\pi} |p(s)| ds \\ &\leq 2\pi K + \int_0^{2\pi} |g(x(s - \tau(s)))| ds + 2\pi \max_{0 \leq s \leq 2\pi} |p(s)|. \end{aligned} \quad (3)$$

We assert that

$$\int_0^{2\pi} |g(x(s - \tau(s)))| ds \leq 2\pi K + 4\pi D_1 \quad (4)$$

for some positive number  $D_1$ . Indeed, integrating equation (1) from 0 to  $2\pi$ , and noting condition (i), we see that

$$\begin{aligned} \int_0^{2\pi} \{g(x(t - \tau(t))) - K\} dt &\leq \int_0^{2\pi} \{g(x(t - \tau(t))) - |f(x'(t))|\} dt \\ &\leq \int_0^{2\pi} \{f(x'(t)) + g(x(t - \tau(t)))\} dt = 0. \end{aligned} \quad (5)$$

Thus letting

$$E_1 = \{t \in [0, 2\pi] \mid x(t - \tau(t)) > D\}, \quad E_2 = [0, 2\pi] \setminus E_1,$$

we have

$$\int_{E_2} |g(x(t - \tau(t)))| dt \leq 2\pi \max \left\{ M, \sup_{|x| \leq D} |g(x)| \right\},$$

and

$$\begin{aligned} \int_{E_1} \{|g(x(t - \tau(t)))| - K\} dt &\leq \int_{E_1} |g(x(t - \tau(t))) - K| dt \\ &= \int_{E_1} \{g(x(t - \tau(t))) - K\} dt \leq - \int_{E_2} \{g(x(t - \tau(t))) - K\} dt \\ &\leq \int_{E_2} |g(x(t - \tau(t)))| dt + \int_{E_2} K dt. \end{aligned}$$

Therefore,

$$\int_0^{2\pi} |g(x(t - \tau(t)))| dt \leq 2\pi K + 4\pi \max \left\{ M, \sup_{|x| \leq D} |g(x)| \right\}$$

as required. Combining (3) and (4), we see that

$$|x'(t)| \leq D_2, \quad t \in [0, 2\pi] \quad (6)$$

for some positive number  $D_2$ . Next, note that the last equality in (5) implies

$$f(x'(t_1)) + g(x(t_1 - \tau(t_1))) = 0$$

for some  $t_1$  in  $[0, 2\pi]$ . Thus in view of condition (i),

$$|g(x(t_1 - \tau(t_1)))| = |f(x'(t_1))| \leq K,$$

and in view of (ii),

$$|x(t_1 - \tau(t_1))| < D.$$

Since  $x(t)$  is  $2\pi$ -periodic, we may infer that  $|x(t_2)| < D$  for some  $t_2$  in  $[0, 2\pi]$ . Finally, we see that

$$\begin{aligned} |x(t)| &= \left| x(t_2) + \int_{t_2}^t x'(t) dt \right| \leq D + \int_0^{2\pi} |x'(t)| dt \\ &\leq D + 2\pi D_2, \quad t \in [0, 2\pi]. \end{aligned} \quad (7)$$

The proof is complete.

By means of the *a priori* bounds just obtained, we may follow the standard procedures as explained in various places of [1] and the continuation theorem in [1, p. 40] to show the existence of a periodic solution of (2). For the sake of completeness, a brief sketch is included as follows.

Let  $X$  be the Banach space of all continuous differentiable functions of the form  $x = x(t)$ , defined on  $R$  such that  $x(t + 2\pi) = x(t)$  for all  $t$ , and endowed with the norm  $\|x\|_1 = \max_{0 \leq t \leq 2\pi} \{|x(t)|, |x'(t)|\}$ . Also let  $Y$  be the Banach space of all continuous functions of the form  $y = y(t)$ , defined on  $R$  such that  $y(t + 2\pi) = y(t)$  for all  $t$ , and endowed with the norm  $\|y\|_0 = \max_{0 \leq t \leq 2\pi} |y(t)|$ . Finally let  $\Omega$  be the subspace of  $X$  containing functions of the form  $x = x(t)$ , such that  $|x(t)| < \bar{D}$  and  $|x'(t)| < \bar{D}$ , where  $\bar{D}$  is a fixed number greater than  $D + 2\pi D_2$ . Now let  $L : X \cap C^{(2)}(R, R) \rightarrow Y$  be the differential operator defined by  $(Lx)(t) = x''(t)$  for  $t \in R$ , and let  $N : X \rightarrow Y$  be defined by

$$(Nx)(t) = -f(x'(t)) - g(x(t - \tau(t))) + p(t), \quad t \in R.$$

Let  $\text{Im } L$  and  $\text{Ker } L$  be, respectively, the image and kernel of the operator  $L$ . Clearly,  $\text{Ker } L = R$ . Furthermore, if we define the projections  $P : X \rightarrow \text{Ker } L$  and  $Q : Y \rightarrow Y/\text{Im } L$  by

$$(Px)(t) = \frac{1}{2\pi} \int_0^{2\pi} x(t) dt, \quad t \in R$$

and

$$(Qy)(t) = \frac{1}{2\pi} \int_0^{2\pi} y(t) dt, \quad t \in R,$$

respectively, then  $\text{Ker } L = \text{Im } P$  and  $\text{Ker } Q = \text{Im } L$ . Furthermore, the operator  $L$  is a Fredholm operator with index zero, and the operator  $N$  is  $L$ -compact on the closure  $\bar{\Omega}$  of  $\Omega$  (see, e.g., [1, p. 176]).

For any  $\lambda \in (0, 1)$  and any  $x = x(t)$  in the domain of  $L$  which also belongs to  $\partial\Omega$ , we must have  $Lx \neq \lambda Nx$ . For otherwise in view of (6) and (7), we see that  $x$  belongs to the interior of  $\Omega$ , which is contrary to the assumption that  $x \in \partial\Omega$ . Next, note that a function  $x = x(t)$  in the intersection of  $\text{Ker } L$  and  $\partial\Omega$  must be the constant functions  $x(t) \equiv \bar{D}$  or  $x(t) \equiv -\bar{D}$ . Hence

$$(QN)(x) = \frac{1}{2\pi} \int_0^{2\pi} (-f(x'(t)) - g(x(t - \tau(t))) + p(t)) dt = -g(\pm \bar{D}) \neq 0.$$

Finally, consider the mapping

$$H(x, s) = sx + (1 - s)g(x), \quad 0 \leq s \leq 1.$$

Since for every  $s \in [0, 1]$  and  $x$  in the intersection of  $\text{Ker } L$  and  $\partial\Omega$ , we have

$$xH(x, s) = sx^2 + (1 - s)xg(x) > 0,$$

thus  $H(x, s)$  is a homotopy. This shows that

$$\begin{aligned} \deg \{QNx, \Omega \cap \text{Ker } L, 0\} &= \deg \{-g(x), \Omega \cap \text{Ker } L, 0\} \\ &= \deg \{-x, \Omega \cap \text{Ker } L, 0\} = \deg \{-x, \Omega \cap R, 0\} \neq 0. \end{aligned}$$

We have thus verified all the assumptions of the continuation theorem [1, p. 40]. Under the assumptions in Theorem 1, equation (2) thus has a  $2\pi$ -periodic solution.

As an example, consider the equation

$$x''(t) + \exp \left\{ - (x'(t))^2 \right\} + \arctan(x(t - \pi)) = \sin t + 1.$$

Take  $f(u) = \exp\{-u^2\} - 1$ ,  $g(u) = \arctan u$  and  $p(u) = \sin u$ , and  $\tau = \pi$ . It is then easy to verify that all the assumptions in Theorem 1 are satisfied with  $K = 1$ ,  $D > \pi/4$ , and  $M = \pi/2$ . Thus this equation has a  $2\pi$ -periodic solution.

## REFERENCES

1. R.E. Gaines and J.L. Mawhin, Coincidence degree and nonlinear differential equations, *Lecture Notes in Math.*, No. 568, Springer-Verlag, (1977).
2. K. Deimling, *Nonlinear Functional Analysis*, Springer-Verlag, (1985).
3. F. Zanolin, Periodic solutions for differential systems of Rayleigh type, *Rend. Istit. Mat. Univ. Trieste* **12** (1/2), 69–77 (1980).
4. S. Invernizzi and F. Zanolin, Periodic solutions of a differential delay equation of Rayleigh type, *Rend. Sem. Mat. Univ. Padova* **61**, 115–124 (1979).
5. F. Liu, Existence of periodic solutions to a class of second-order nonlinear differential equations, *Acta Math. Sinica* **33** (2), 260–269 (1990).
6. P. Omari and G. Villari, Periodic solutions of the Rayleigh equations with damping of definite sign, *Atti. Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur.* **1** (1), 29–35 (1990).